

The dinner table problem: the rectangular case

Roberto Tauraso *


Dipartimento di Matematica
Università di Roma “Tor Vergata”
00133 Roma, Italy
tauraso@mat.uniroma2.it

February 2, 2008

1 Introduction

Assume that 8 people are seated around a table and we want to enumerate the number of ways that they can be permuted such that neighbors are no more neighbors after the rearrangement. Of course the answer depends on the *topology* of the table: if the table is a circle then it is easy to check by a simple computer program that the permutations that verify this property are 2832, if it is a long bar and all the persons seat along one side then they are 5242. Furthermore, if it is a rectangular table with two sides then the rearrangements are 9512. The first two cases are respectively described by sequences A089222 and A002464 of the On-Line Encyclopedia of Integer Sequences [3], on the other hand the rectangular case does not appear in the literature.

Here is a valid rearrangement for $n = 8$:



whose associated permutation is


$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 2 & 6 & 8 & 3 & 4 & 7 & 5 \end{pmatrix}.$$

*I would like to warmly thank Alessandro Nicolosi and Giorgio Minenkov for drawing my attention to this problem.

For a generic number of persons n the required property can be established more formally in this way:

$$|\pi(i+2) - \pi(i)| \neq 2 \quad \text{for } 1 \leq i \leq n-2.$$

It is interesting to note that this rearrangement problem around a table has also another remarkable interpretation. Consider n kings to be placed on a $n \times n$ board, one in each row and column, in such a way that they are non-attacking with respect to these different *topologies* of the board: if we enumerate the ways on a toroidal board we find the sequence *A089222*, for a regular board we have *A002464*, and finally if we divide the board in the main four quadrants we are considering the new sequence (now recorded as *A110128*). Here is the 8 kings displacement that corresponds to the permutation π introduced before:



Closed formulas for the first two sequences are known: for *A089222* it is (see [1])

$$a_{n,0} = \sum_{r=0}^{n-1} \sum_{c=0}^r (-1)^r 2^c (n-r)! \binom{r-1}{c-1} \binom{n-r}{c} \left(\frac{n}{n-r} \right)^2 + (-1)^n 2n$$

and for *A002464* it is (see [2])

$$a_{n,1} = \sum_{r=0}^{n-1} \sum_{c=0}^r (-1)^r 2^c (n-r)! \binom{r-1}{c-1} \binom{n-r}{c}$$

(remember that $\binom{k}{-1} = 0$ if $k \neq -1$ and $\binom{-1}{-1} = 1$).

In this paper we propose to study the sequence $a_{n,d}$ defined for $2 \leq d \leq n-1$ as follows: $a_{n,d}$ denotes the total number of permutations π of $\{1, 2, \dots, n\}$ such that

$$|\pi(i+d) - \pi(i)| \neq d \quad \text{for } 1 \leq i \leq n-d.$$

The table below provides some numerical values:

n	$a_{n,0}$	$a_{n,1}$	$a_{n,2}$	$a_{n,3}$
1	1	1	1	1
2	0	0	2	2
3	0	0	4	6
4	0	2	16	20
5	10	14	44	80
6	36	90	200	384
7	322	646	1288	2240
8	2832	5242	9512	15424
9	27954	47622	78652	123456
10	299260	479306	744360	1110928
11	3474482	5296790	7867148	11287232
12	43546872	63779034	91310696	127016304
13	586722162	831283558	1154292796	1565107248
14	8463487844	11661506218	15784573160	20935873872
15	130214368530	175203184374	232050062524	301974271248
16	2129319003680	2806878055610	3648471927912	4669727780624

We will show that the following formula holds for $d \geq 2$:

$$a_{n,d} = \sum_{r_1=0}^{n_1-1} \sum_{c_1=0}^{r_0} \cdots \sum_{r_d=0}^{n_d-1} \sum_{c_d=0}^{r_d} (-1)^r 2^c (n-r-c)! \prod_{k=1}^d \binom{n_k - r_k}{c_k} \sum_{\substack{\sum_{i=1}^{c_1} l_{i,1}=r_1 \\ l_{i,1} \geq 1}} \cdots \sum_{\substack{\sum_{i=1}^{c_d} l_{i,d}=r_d \\ l_{i,d} \geq 1}} q_{n,d}(L)$$

where $n_k = |N_k| = |\{1 \leq i \leq n : i \equiv k \pmod{d}\}|$, $r = \sum_{k=1}^d r_k$, $c = \sum_{k=1}^d c_k$ and

$$q_{n,d}(L) = q_{n,d}([l_1, \dots, l_c]) = \sum_{\substack{J_1 \dot{\cup} \dots \dot{\cup} J_d = \{1, \dots, c\} \\ n_k \geq \sum_{i \in J_k} l_i}} \prod_{k=1}^d \binom{n_k - \sum_{i \in J_k} l_i}{|J_k|} \cdot |J_k|!.$$

In the last section, we will prove that the probability that a permutation belongs to the set enumerated by $a_{n,d}$ tends always to e^{-2} as n goes to infinity. A more precise expansion reveals how the limit probability depends on d :

$$\frac{a_{n,d}}{n!} = e^{-2} \left(1 + \frac{4(d-1)}{n} + O\left(\frac{1}{n^2}\right) \right).$$

2 Asymptotic analysis: cases $d = 0$ and $d = 1$

Proposition 2.1. *The following asymptotic expansions hold:*

$$\frac{a_{n,0}}{n!} \sim e^{-2} \left(1 - \frac{4}{n} + \frac{20}{3n^3} + \frac{58}{3n^4} + \frac{736}{15n^5} + O\left(\frac{1}{n^6}\right) \right),$$

and

$$\frac{a_{n,1}}{n!} \sim e^{-2} \left(1 - \frac{2}{n^2} - \frac{10}{3n^3} - \frac{6}{n^4} - \frac{154}{15n^5} + O\left(\frac{1}{n^6}\right) \right),$$

Proof. The expansion of $a_{n,0}/n!$ is contained in [1] and it was obtained from a recurrence relation by the method of undetermined coefficients. However, since we are particularly interested in, we will explicitly derive the coefficient of $1/n$.

As regards $a_{n,1}/n!$, we give the detailed proof only for the coefficients of $1/n$ and $1/n^2$ (the others can be computed in a similar way). Since

$$\frac{a_{n,1}}{n!} = \sum_{r=0}^{n-1} \sum_{c=0}^r (-1)^r \frac{2^c}{c!} \binom{r-1}{c-1} \frac{n^{r+c}}{(n^x)^2}$$

and as n goes to infinity the resulting series is uniformly convergent (with respect to n), then we can consider each term one by one. Moreover, it suffices to analyse the cases when c is equal to r , $r-1$ and $r-2$ because the rational function $n^{r+c}/(n^x)^2 \sim 1/n^{r-c}$. For $c = r$

$$\binom{r-1}{r-1} \frac{n^{2r}}{(n^x)^2} \sim \frac{1 - \left[\frac{2r}{2r-1} \right] \frac{1}{n} + \left[\frac{2r}{2r-2} \right] \frac{1}{n^2}}{\left(1 - \left[\frac{r}{r-1} \right] \frac{1}{n} + \left[\frac{r}{r-2} \right] \frac{1}{n^2} \right)^2} \sim 1 - \frac{r^2 + r}{n} + \frac{r^4 + 4r^3 + 2r^2}{2n^2}.$$

In a similar way, for $c = r-1$

$$\binom{r-1}{r-2} \frac{n^{2r-1}}{(n^x)^2} \sim \frac{r-1}{n} - \frac{(r-1)^3 + 3(r-1)^2 + (r-1)}{n^2}$$

and for $c = r-2$

$$\binom{r-1}{r-3} \frac{n^{2r-2}}{(n^x)^2} \sim \frac{(r-2)^2 + 2(r-2)}{2n^2}.$$

Hence

$$\begin{aligned} \frac{a_{n,1}}{n!} &\sim \sum_{r=0}^{n-1} \frac{(-2)^r}{r!} \left(1 - \frac{r^2 + r}{n} + \frac{r^4 + 4r^3 + 2r^2}{2n^2} \right) + \\ &\quad - \sum_{r=1}^{n-1} \frac{(-2)^{r-1}}{(r-1)!} \left(\frac{r-1}{n} - \frac{(r-1)^3 + 3(r-1)^2 + (r-1)}{n^2} \right) + \\ &\quad + \sum_{r=2}^{n-1} \frac{(-2)^{r-2}}{(r-2)!} \left(\frac{(r-2)^2 + 2(r-2)}{2n^2} \right). \end{aligned}$$

Taking the sums we find that

$$\begin{aligned} \frac{a_{n,1}}{n!} &\sim e^{-2} \left(1 + \frac{(-2^2 + 2) + 2}{n} + \frac{(2^4 - 4 \cdot 2^3 + 2 \cdot 2^2) + 2 \cdot (-2^3 + 3 \cdot 2^2 - 2) + (2^2 - 2 \cdot 2)}{2n^2} \right) \\ &\sim e^{-2} \left(1 - \frac{2}{n^2} \right). \end{aligned}$$

Note that the same strategy can be applied also to $a_{n,0}/n!$. For example the coefficient of $1/n$ can be easily calculated: for $c = r$ the formula gives

$$\binom{r-1}{r-1} \frac{n^{2r}}{(n^r)^2} \cdot \left(\frac{n}{n-r}\right)^2 \sim \left(1 - \frac{r^2+r}{n}\right) \cdot \left(1 + \frac{2r}{n}\right) \sim 1 - \frac{r^2-r}{n}$$

and therefore

$$\begin{aligned} \frac{a_{n,0}}{n!} &\sim \sum_{r=0}^{n-1} \frac{(-2)^r}{r!} \left(1 - \frac{r^2-r}{n}\right) - \sum_{r=1}^{n-1} \frac{(-2)^{r-1}}{(r-1)!} \left(\frac{r-1}{n}\right) \\ &\sim e^{-2} \left(1 + \frac{(-2^2-2)+2}{n}\right) \sim e^{-2} \left(1 - \frac{4}{n}\right). \end{aligned}$$

□

3 The formula for $d \geq 2$

Theorem 3.1. For $d \geq 2$

$$a_{n,d} = \sum_{r_1=0}^{n_1-1} \sum_{c_1=0}^{r_0} \cdots \sum_{r_d=0}^{n_d-1} \sum_{c_d=0}^{r_d} (-1)^r 2^c (n-r-c)! \prod_{k=1}^d \binom{n_k - r_k}{c_k} \sum_{\substack{\sum_{i=1}^{c_1} l_{i,1}=r_1 \\ l_{i,1} \geq 1}} \cdots \sum_{\substack{\sum_{i=1}^{c_d} l_{i,d}=r_d \\ l_{i,d} \geq 1}} q_{n,d}(L)$$

where $n_k = |N_k| = |\{1 \leq i \leq n : i \equiv k \pmod{d}\}|$, $r = \sum_{k=1}^d r_k$, $c = \sum_{k=1}^d c_k$ and

$$q_{n,d}(L) = q_{n,d}([l_1, \dots, l_c]) = \sum_{\substack{J_1 \dot{\cup} \dots \dot{\cup} J_d = \{1, \dots, c\} \\ n_k \geq \sum_{i \in J_k} l_i}} \prod_{k=1}^d \binom{n_k - \sum_{i \in J_k} l_i}{|J_k|} \cdot |J_k|!.$$

Remark 3.1. Note that for $d = 1$ the above formula coincides with the one due to Robbins:

$$a_{n,1} = \sum_{r=0}^{n-1} \sum_{c=0}^r (-1)^r 2^c (n-r-c)! \binom{n-r}{c} \sum_{\substack{l_1 + \dots + l_c = r \\ l_i \geq 1}} q_{n,1}([l_1, \dots, l_c]),$$

where

$$\begin{aligned} \sum_{\substack{l_1 + \dots + l_c = r \\ l_i \geq 1}} q_{n,1}([l_1, \dots, l_c]) &= \sum_{\substack{l_1 + \dots + l_c = r \\ l_i \geq 1}} \sum_{\substack{J = \{1, \dots, c\} \\ n \geq r}} \binom{n - \sum_{i \in J} l_i}{|J|} (|J|!) \\ &= \sum_{\substack{l_1 + \dots + l_c = r \\ l_i \geq 1}} \binom{n-r}{c} c! = \binom{r-1}{c-1} \binom{n-r}{c} c!. \end{aligned}$$

Proof of the Theorem. For $i = 1, \dots, n-d$, let $T_{i,d}$ be the set of permutations of $\{1, 2, \dots, n\}$ such that i and $i+d$ are d -consecutive:

$$T_{i,d} = \{\pi \in S_n : |\pi^{-1}(i+d) - \pi^{-1}(i)| = d\}$$

(for $i = n-d+1, \dots, n$ we consider $T_{i,d}$ as an empty set). Then by the Inclusion-Exclusion Principle

$$a_{n,d} = \sum_{I \subset \{1,2,\dots,n\}} (-1)^{|I|} \left| \bigcap_{i \in I} T_{i,d} \right|$$

assuming the convention that when the intersection is made over an empty set of indices then it is the whole set of permutations S_n . A d -component of a set of indices I is a maximal subset of d -consecutive integers, and we denote by $\sharp I$ the number of d -components of I . So the above formula can be rewritten as

$$a_{n,d} = \sum_{r_1=0}^{n_1-1} \sum_{c_1=0}^{r_1} \cdots \sum_{r_d=0}^{n_d-1} \sum_{c_d=0}^{r_d} (-1)^r \left| \bigcap_{i \in I_1 \cup \dots \cup I_d} T_{i,d} \right|$$

$I_k \subset N_k, |I_k| = r_k, \sharp I_k = c_k$


where $N_k = \{1 \leq i \leq n : i \equiv k \pmod{d}\}$ and $n_k = |N_k|$.

Remark 3.2. In order to better illustrate the idea of the proof, which is inspired by the one of Robbins in [2], we give an example of how a permutation π which belongs to the above intersection of $T_{i,d}$'s can be selected. Assume that $n = 12$, $d = 2$, $r_1 = 2$, $c_1 = 1$, $r_2 = 3$, $c_2 = 2$. N_1 and N_2 are respectively the odd numbers and the even numbers between 1 and 12. Now we choose I_1 and I_2 : let $l_{i,k}$ be the component sizes (each component fixes actually $l_{i,k} + 1$ numbers) and let $j_{i,k}$ be the sizes of the gaps between consecutive components. Then the choice of I_1 and I_2 is equivalent to select an integral solution of

$$\begin{cases} l_{1,1} = 2 & , \quad l_{i,1} \geq 1 \\ l_{1,2} + l_{2,2} = 3 & , \quad l_{i,2} \geq 1 \\ j_{0,1} + j_{1,1} = 3 & , \quad j_{i,1} \geq 0 \\ j_{0,2} + j_{1,2} + j_{2,2} = 1 & , \quad j_{i,2} \geq 0 \end{cases}$$

For example taking $l_{1,1} = 2$, $l_{1,2} = 1$, $l_{2,2} = 2$, $j_{0,1} = 1$, $j_{1,1} = 2$, $j_{0,2} = 0$, $j_{1,2} = 1$ and $j_{2,2} = 0$, we select the set of indices $I_1 = \{3, 5\}$ and $I_2 = \{2, 8, 10\}$.


Here is the corresponding table arrangement:



Now we redistribute the three components selecting a partition $J_1 \overset{\circ}{\cup} J_2 = \{1, 2, 3\}$, say $J_1 = \{1, 2\}$ and $J_2 = \{3\}$. This means that the first two components $\{2\}$ and $\{3, 5\}$ will go to the odd seats and the third component $\{8, 10\}$ will go to the even seats. Then we decide the component displacements and orientations: for example in the odd seats we place first $\{3, 5\}$ reversed and then $\{2\}$ and in the even seats we place $\{8, 10\}$ reversed. To determine the component positions we need the new gap sizes and therefore we solve the following two equations

$$\begin{cases} j'_{0,1} + j'_{1,1} + j'_{2,1} = 1 & , \quad j'_{i,1} \geq 0 \\ j'_{0,2} + j'_{1,2} = 3 & , \quad j'_{i,2} \geq 0 \end{cases}.$$

If we take $j'_{0,1} = 0$, $j'_{1,1} = 1$, $j'_{2,1} = 0$, $j'_{0,2} = 1$, $j'_{1,2} = 2$ and we fill the empty places with the remaining numbers 1, 6, 9 and 11, we obtain the following table rearrangement:



We go back to the proof. Each I_k is determined by an integral solution of

$$\begin{cases} l_{1,k} + l_{2,k} + \dots + l_{c_k,k} = r_k & , \quad l_{i,k} \geq 1 \\ j_{0,k} + j_{1,k} + \dots + j_{c_k,k} = n_k - r_k - c_k & , \quad j_{i,k} \geq 0 \end{cases}$$

where $l_{1,k}, \dots, l_{c_k,k}$ are the sizes of each component and $j_{0,k}, \dots, j_{c_k,k}$ are the sizes of the gaps. The counting of the number of integral solutions of this systems yields the following factor in the formula

$$\prod_{k=1}^d \binom{n_k - r_k}{c_k} \sum_{\substack{\sum_{i=1}^{c_1} l_{i,1} = r_1 \\ l_{i,1} \geq 1}} \dots \sum_{\substack{\sum_{i=1}^{c_d} l_{i,d} = r_d \\ l_{i,d} \geq 1}}.$$

Now, given $I_1, \dots, I_d$, we select a permutation $\pi \in |\bigcap_{i \in I_1 \cup \dots \cup I_d} T_{i,d}|$:

- we redistribute the $c = \sum_{k=1}^d c_k$ components selecting a partition $J_1, \dots, J_d$ of the set $\{1, \dots, c\}$ (we allow that J_k can be empty),
- for each set J_k of the partition, we determine the sizes of the new gaps solving $j'_{0,k} + j'_{1,k} + \dots + j'_{|J_k|,k} = n_k - \sum_{i \in J_k} l_i - |J_k|$, $j'_{i,k} \geq 0$,
- we choose the order of components and their orientation in $|J_k|! \cdot 2^{|J_k|}$ ways

- we fill the empty places with the remaining numbers in $(n - r - c)!$ ways.

Taking into account all these effects we obtain

$$\sum_{\substack{J_1 \dot{\cup} \dots \dot{\cup} J_d = \{1, \dots, c\} \\ n_k \geq \sum_{i \in J_k} l_i}} (n - r - c)! \prod_{k=1}^d \binom{n_k - \sum_{i \in J_k} l_i}{|J_k|} \cdot |J_k|! \cdot 2^{|J_k|}.$$

Finally, since $c = \sum_{k=1}^d |J_k|$, it is easy to get the required formula. $\square$

4 Asymptotic analysis: the general case

Theorem 4.1. *For $d \geq 0$*

$$\frac{a_{n,d}}{n!} = e^{-2} \left(1 + \frac{4(d-1)}{n} + O\left(\frac{1}{n^2}\right) \right).$$

Proof. The cases $d = 0$ and $d = 1$ have been already discussed, so we assume that $d \geq 2$. By the general formula

$$\frac{a_{n,d}}{n!} = \sum \frac{\text{polynomial in } n \text{ of degree } 2c}{\text{polynomial in } n \text{ of degree } r + c}.$$

Since $c_k \leq r_k$, the rational functions go to zero faster than $1/n$ as n goes to infinity, unless either $r + c = 2c$ or $r + c = 2c + 1$. In the first case $c_k = r_k$ and $l_{i,k} = 1$, in the second case there is one and only one index k such that $c_k + 1 = r_k$ and $l_{i,k} = 2$ for some i . By symmetry, we can assume that that particular index $k = d$ and multiply the corresponding term by d :

$$\begin{aligned} \frac{a_{n,d}}{n!} \sim & \sum_{r_1=0}^{n/d-1} \dots \sum_{r_d=0}^{n/d-1} \left((-2)^r \frac{(n-2r)!}{n!} \prod_{k=1}^d \binom{n/d - r_k}{r_k} q_{n,d}(\overbrace{[1, \dots, 1]}^r) + \right. \\ & \left. + d \cdot (-2)^{r-1} \frac{(n-2r+1)!}{n!} \prod_{k=1}^d \binom{n/d - r_k}{r_k} \binom{n/d - r_d}{r_d - 1} (r_d - 1) q_{n,d}(\overbrace{[1, \dots, 1, 2]}^{r-2}) \right), \end{aligned}$$

that is

$$\begin{aligned} \frac{a_{n,d}}{n!} \sim & \sum_{r_1=0}^{n/d-1} \dots \sum_{r_d=0}^{n/d-1} \frac{q_{n,d}(\overbrace{[1, \dots, 1]}^r)}{n^{2r}} \prod_{k=1}^d \frac{(-2)^{r_k}}{r_k!} \prod_{k=1}^d \frac{(n/d)^{2r_k}}{(n/d)^{r_k}} + \\ & + 2d \cdot \sum_{r_1=0}^{n/d-1} \dots \sum_{r_{d-1}=0}^{n/d-1} \sum_{r_d=2}^{n/d-1} \frac{q_{n,d}(\overbrace{[1, \dots, 1, 2]}^{r-2})}{n^{2r-1}} \prod_{k=1}^{d-1} \frac{(-2)^{r_k}}{r_k!} \cdot \frac{(-2)^{r_d-2}}{(r_d-2)!} \prod_{k=1}^{d-1} \frac{(n/d)^{2r_k}}{(n/d)^{r_k}} \cdot \frac{(n/d)^{2r_d-1}}{(n/d)^{r_d}}. \end{aligned}$$

We start considering the second term. Since

$$\begin{aligned}
q_{n,d}(\overbrace{[1, \dots, 1]}^{r-2}, 2] &\sim \sum_{\substack{\sum_{k=1}^d r'_k = r-1 \\ r'_k \geq 0}} \binom{r-1}{r'_1, \dots, r'_d} \prod_{k=1}^{d-1} \binom{n/d - r'_k}{r'_k} \cdot r'_k! \cdot \binom{n/d - r'_d - 1}{r'_d} \cdot r'_d! \\
&\sim \sum_{\substack{\sum_{k=1}^d r'_k = r-1 \\ r'_k \geq 0}} \binom{r-1}{r'_1, \dots, r'_d} \prod_{k=1}^{d-1} \frac{(n/d)^{\underline{2r'_k}}}{(n/d)^{\underline{r'_k}}} \cdot \frac{(n/d)^{\underline{2r'_d+1}}}{(n/d)^{\underline{r'_d+1}}} \\
&\sim \left(\frac{n}{d}\right)^{r-1} \sum_{\substack{\sum_{k=1}^d r'_k = r-1 \\ r'_k \geq 0}} \binom{r-1}{r'_1, \dots, r'_d} = \left(\frac{n}{d}\right)^{r-1} d^{r-1} = n^{r-1}
\end{aligned}$$

then it becomes

$$2 \cdot \sum_{r_1=0}^{n/d-1} \cdots \sum_{r_{d-1}=0}^{n/d-1} \sum_{r_d=2}^{n/d-1} \frac{n^{r-1}}{n^{2r-1}} \prod_{k=1}^{d-1} \frac{(-2/d)^{r_k}}{r_k!} \cdot \frac{(-2/d)^{r_d-2}}{(r_d-2)!} \cdot n^{r-1} \sim 2 (e^{-2/d})^d \frac{1}{n} = \frac{2e^{-2}}{n}.$$

Now we consider the first term. Since

$$\begin{aligned}
q_{n,d}(\overbrace{[1, \dots, 1]}^r) &\sim \sum_{\substack{\sum_{k=1}^d r'_k = r \\ r'_k \geq 0}} \binom{r}{r'_1, \dots, r'_d} \prod_{k=1}^d \binom{n/d - r'_k}{r'_k} \cdot r'_k! \\
&\sim \sum_{\substack{\sum_{k=1}^d r'_k = r \\ r'_k \geq 0}} \binom{r}{r'_1, \dots, r'_d} \prod_{k=1}^d \frac{(n/d)^{\underline{2r'_k}}}{(n/d)^{\underline{r'_k}}} \\
&\sim \left(\frac{n}{d}\right)^r \sum_{\substack{\sum_{k=1}^d r'_k = r \\ r'_k \geq 0}} \binom{r}{r'_1, \dots, r'_d} \prod_{k=1}^d \frac{1 - \frac{1}{2}(2r'_k)(2r'_k - 1) \cdot \frac{d}{n}}{1 - \frac{1}{2}r'_k(r'_k - 1) \cdot \frac{d}{n}} \\
&\sim \left(\frac{n}{d}\right)^r \sum_{\substack{\sum_{k=1}^d r'_k = r \\ r'_k \geq 0}} \binom{r}{r'_1, \dots, r'_d} \left(1 - \left(\frac{3}{2} \sum_{k=1}^d r'_k(r'_k - 1) + \sum_{k=1}^d r'_k\right) \cdot \frac{d}{n}\right) \\
&\sim \left(\frac{n}{d}\right)^r \left(d^r - \left(\frac{3}{2}dr(r-1)d^{r-2} + drd^{r-1}\right) \cdot \frac{d}{n}\right) \\
&\sim n^r \left(1 - \left(\frac{3}{2}r(r-1) + dr\right) \frac{1}{n}\right)
\end{aligned}$$

then it becomes

$$\sum_{r_1=0}^{n/d-1} \cdots \sum_{r_d=0}^{n/d-1} \frac{1 - \left(\frac{3}{2}r(r-1) + dr\right) \cdot \frac{1}{n}}{1 - r(2r-1) \cdot \frac{1}{n}} \prod_{k=1}^d \frac{(-2/d)^{r_k}}{r_k!} \prod_{k=1}^d \left(\frac{1 - r_k(2r_k-1) \cdot \frac{d}{n}}{1 - \frac{1}{2}r_k(r_k-1) \cdot \frac{d}{n}} \right)$$

that is

$$\sum_{r_1=0}^{n/d-1} \cdots \sum_{r_d=0}^{n/d-1} \left(1 + \left(\frac{r^2 + r}{2} - dr \right) \cdot \frac{1}{n} \right) \prod_{k=1}^d \frac{(-2/d)^{r_k}}{r_k!} \prod_{k=1}^d \left(1 - \left(\frac{3r_k(r_k-1)}{2} + r_k \right) \cdot \frac{d}{n} \right).$$

Reminding that $r = \sum_{k=1}^d r_k$, it is equivalent to

$$\begin{aligned} e^{-2} + \frac{1}{n} \cdot \sum_{r_1=0}^{n/d-1} \cdots \sum_{r_d=0}^{n/d-1} & \left(\frac{1-3d}{2} \sum_{k=1}^d r_k(r_k-1) \prod_{k=1}^d \frac{(-2/d)^{r_k}}{r_k!} + \right. \\ & \left. + (1-2d) \sum_{k=1}^d r_k \prod_{k=1}^d \frac{(-2/d)^{r_k}}{r_k!} + \sum_{k=1}^d \sum_{k'=k+1}^d r_k r_{k'} \prod_{k=1}^d \frac{(-2/d)^{r_k}}{r_k!} \right) \end{aligned}$$

and taking the limit of the remaining sums we obtain

$$e^{-2} \left(1 + \frac{1}{n} \cdot \left(\frac{1-3d}{2} \left(\frac{-2}{d} \right)^2 d + (1-2d) \left(\frac{-2}{d} \right) d + \left(\frac{-2}{d} \right)^2 \frac{d(d-1)}{2} \right) \right)$$

that is

$$e^{-2} \left(1 + \frac{4d-6}{n} \right).$$

Finally, putting all together we find that

$$\frac{a_{n,d}}{n!} \sim e^{-2} \left(1 + \frac{4d-6}{n} \right) + \frac{2e^{-2}}{n} = e^{-2} \left(1 + \frac{4(d-1)}{n} \right).$$

□

References

- [1] B. Aspvall and F. M. Liang. The dinner table problem. Technical Report STAN-CS-80-829, Computer Science Department, Stanford University, Stanford, California, 1980.
- [2] D. P. Robbins. The probability that neighbors remain neighbors after random rearrangements, *Amer. Math. Monthly*, **87** (1980), 122–124.
- [3] N. J. A. Sloane. The On-Line Encyclopedia of Integer Sequences, on the web at <http://www.research.att.com/~njas/sequences/>.